

MATH 311

Topics in Applied Mathematics I


Lecture 21b:
Eigenvalues and eigenvectors.
Characteristic equation.

Linear transformations of \mathbb{R}^2


Any linear mapping $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is represented as multiplication of a 2-dimensional column vector by a 2×2 matrix: $f(\mathbf{x}) = A\mathbf{x}$ or

$$f \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$


Linear transformations corresponding to particular matrices can have various geometric properties.




$$A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$




Rotation by 90°




$$A = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$$




Rotation by 45°




$$A = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$$




Reflection about
the vertical axis




$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$




Reflection about
the line $x - y = 0$




$$A = \begin{pmatrix} 1 & 1/2 \\ 0 & 1 \end{pmatrix}$$




Horizontal shear




$$A = \begin{pmatrix} 1/2 & 0 \\ 0 & 1/2 \end{pmatrix}$$




Scaling




$$A = \begin{pmatrix} 3 & 0 \\ 0 & 1/3 \end{pmatrix}$$




Squeeze




$$A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$




Vertical projection on
the horizontal axis




$$A = \begin{pmatrix} 0 & -1 \\ 0 & 1 \end{pmatrix}$$



Horizontal projection
on the line $x + y = 0$



$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$



Identity

Eigenvalues and eigenvectors

Definition. Let A be an $n \times n$ matrix. A number $\lambda \in \mathbb{R}$ is called an **eigenvalue** of the matrix A if $A\mathbf{v} = \lambda\mathbf{v}$ for a nonzero column vector $\mathbf{v} \in \mathbb{R}^n$.

The vector \mathbf{v} is called an **eigenvector** of A belonging to (or associated with) the eigenvalue λ .

Remarks.

- Alternative notation:
eigenvalue = **characteristic value**,
eigenvector = **characteristic vector**.

- The zero vector is never considered an eigenvector.

Example. $A = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$.

$$\begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

$$\begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix} \begin{pmatrix} 0 \\ -2 \end{pmatrix} = \begin{pmatrix} 0 \\ -6 \end{pmatrix} = 3 \begin{pmatrix} 0 \\ -2 \end{pmatrix}.$$

Hence $(1, 0)$ is an eigenvector of A belonging to the eigenvalue 2, while $(0, -2)$ is an eigenvector of A belonging to the eigenvalue 3.

Example. $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

Hence $(1, 1)$ is an eigenvector of A belonging to the eigenvalue 1, while $(1, -1)$ is an eigenvector of A belonging to the eigenvalue -1 .

Vectors $\mathbf{v}_1 = (1, 1)$ and $\mathbf{v}_2 = (1, -1)$ form a basis for \mathbb{R}^2 . Consider a linear operator $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $L(\mathbf{x}) = A\mathbf{x}$. The matrix of L with respect to the basis $\mathbf{v}_1, \mathbf{v}_2$ is $B = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Let A be an $n \times n$ matrix. Consider a linear operator $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by $L(\mathbf{x}) = A\mathbf{x}$.

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ be a nonstandard basis for \mathbb{R}^n and B be the matrix of the operator L with respect to this basis.

Theorem The matrix B is diagonal if and only if vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are eigenvectors of A .

If this is the case, then the diagonal entries of the matrix B are the corresponding eigenvalues of A .

$$A\mathbf{v}_i = \lambda_i\mathbf{v}_i \iff B = \begin{pmatrix} \lambda_1 & & & 0 \\ & \lambda_2 & & \\ & & \ddots & \\ 0 & & & \lambda_n \end{pmatrix}$$

Eigenspaces

Let A be an $n \times n$ matrix. Let \mathbf{v} be an eigenvector of A belonging to an eigenvalue λ .

Then $A\mathbf{v} = \lambda\mathbf{v} \implies A\mathbf{v} = (\lambda I)\mathbf{v} \implies (A - \lambda I)\mathbf{v} = \mathbf{0}$.

Hence $\mathbf{v} \in N(A - \lambda I)$, the nullspace of the matrix $A - \lambda I$.

Conversely, if $\mathbf{x} \in N(A - \lambda I)$ then $A\mathbf{x} = \lambda\mathbf{x}$.

Thus the eigenvectors of A belonging to the eigenvalue λ are nonzero vectors from $N(A - \lambda I)$.

Definition. If $N(A - \lambda I) \neq \{\mathbf{0}\}$ then it is called the **eigenspace** of the matrix A corresponding to the eigenvalue λ .

How to find eigenvalues and eigenvectors?

Theorem Given a square matrix A and a scalar λ , the following statements are equivalent:

- λ is an eigenvalue of A ,
- $N(A - \lambda I) \neq \{\mathbf{0}\}$,
- the matrix $A - \lambda I$ is singular,
- $\det(A - \lambda I) = 0$.

Definition. $\det(A - \lambda I) = 0$ is called the **characteristic equation** of the matrix A .

Eigenvalues λ of A are roots of the characteristic equation. Associated eigenvectors of A are nonzero solutions of the equation $(A - \lambda I)\mathbf{x} = \mathbf{0}$.

Example. $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$

$$\begin{aligned} \det(A - \lambda I) &= \begin{vmatrix} a - \lambda & b \\ c & d - \lambda \end{vmatrix} \\ &= (a - \lambda)(d - \lambda) - bc \\ &= \lambda^2 - (a + d)\lambda + (ad - bc). \end{aligned}$$

Example. $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}.$

$$\det(A - \lambda I) = \begin{vmatrix} a_{11} - \lambda & a_{12} & a_{13} \\ a_{21} & a_{22} - \lambda & a_{23} \\ a_{31} & a_{32} & a_{33} - \lambda \end{vmatrix}$$
$$= -\lambda^3 + c_1\lambda^2 - c_2\lambda + c_3,$$

where $c_1 = a_{11} + a_{22} + a_{33}$ (the *trace* of A),

$$c_2 = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} + \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} + \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix},$$

$$c_3 = \det A.$$

Theorem. Let $A = (a_{ij})$ be an $n \times n$ matrix.

Then $\det(A - \lambda I)$ is a polynomial of λ of degree n :

$$\det(A - \lambda I) = (-1)^n \lambda^n + c_1 \lambda^{n-1} + \cdots + c_{n-1} \lambda + c_n.$$

Furthermore, $(-1)^{n-1} c_1 = a_{11} + a_{22} + \cdots + a_{nn}$
and $c_n = \det A$.

Definition. The polynomial $p(\lambda) = \det(A - \lambda I)$ is called the **characteristic polynomial** of the matrix A .

Corollary Any $n \times n$ matrix has at most n eigenvalues.