

MATH 323  
Linear Algebra

**Lecture 11:**  
**Linear independence (continued).**  
**Basis and dimension.**

## Linear independence

*Definition.* Let  $V$  be a vector space. Vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in V$  are called **linearly dependent** if they satisfy a relation

$$r_1\mathbf{v}_1 + r_2\mathbf{v}_2 + \dots + r_k\mathbf{v}_k = \mathbf{0},$$

where the coefficients  $r_1, \dots, r_k \in \mathbb{R}$  are not all equal to zero. Otherwise vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  are called **linearly independent**. That is, if

$$r_1\mathbf{v}_1 + r_2\mathbf{v}_2 + \dots + r_k\mathbf{v}_k = \mathbf{0} \implies r_1 = \dots = r_k = 0.$$

A set  $S \subset V$  is **linearly dependent** if one can find some distinct linearly dependent vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k$  in  $S$ . Otherwise  $S$  is **linearly independent**.

**Theorem** The following conditions are equivalent:

(i) vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k$  are linearly dependent;

(ii) one of vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k$  is a linear combination of the other  $k - 1$  vectors.

*Proof:* (i)  $\implies$  (ii) Suppose that

$$r_1\mathbf{v}_1 + r_2\mathbf{v}_2 + \cdots + r_k\mathbf{v}_k = \mathbf{0},$$

where  $r_i \neq 0$  for some  $1 \leq i \leq k$ . Then

$$\mathbf{v}_i = -\frac{r_1}{r_i}\mathbf{v}_1 - \cdots - \frac{r_{i-1}}{r_i}\mathbf{v}_{i-1} - \frac{r_{i+1}}{r_i}\mathbf{v}_{i+1} - \cdots - \frac{r_k}{r_i}\mathbf{v}_k.$$

(ii)  $\implies$  (i) Suppose that

$$\mathbf{v}_i = s_1\mathbf{v}_1 + \cdots + s_{i-1}\mathbf{v}_{i-1} + s_{i+1}\mathbf{v}_{i+1} + \cdots + s_k\mathbf{v}_k$$

for some scalars  $s_j$ . Then

$$s_1\mathbf{v}_1 + \cdots + s_{i-1}\mathbf{v}_{i-1} - \mathbf{v}_i + s_{i+1}\mathbf{v}_{i+1} + \cdots + s_k\mathbf{v}_k = \mathbf{0}.$$

**Problem.** Let  $A = \begin{pmatrix} -1 & 1 \\ -1 & 0 \end{pmatrix}$ . Determine whether matrices  $A$ ,  $A^2$ , and  $A^3$  are linearly independent.

$$\text{We have } A = \begin{pmatrix} -1 & 1 \\ -1 & 0 \end{pmatrix}, \quad A^2 = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}, \quad A^3 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

The task is to check if there exist  $r_1, r_2, r_3 \in \mathbb{R}$  not all zero such that  $r_1A + r_2A^2 + r_3A^3 = O$ .

This matrix equation is equivalent to a system

$$\begin{cases} -r_1 + 0r_2 + r_3 = 0 \\ r_1 - r_2 + 0r_3 = 0 \\ -r_1 + r_2 + 0r_3 = 0 \\ 0r_1 - r_2 + r_3 = 0 \end{cases} \quad \left( \begin{array}{ccc|c} -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \end{array} \right) \rightarrow \left( \begin{array}{ccc|c} 1 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

The row echelon form of the augmented matrix shows there is a free variable. Hence the system has a nonzero solution so that the matrices are linearly dependent (one relation is  $A + A^2 + A^3 = O$ ).

**Problem.** Show that functions  $e^x$ ,  $e^{2x}$ , and  $e^{3x}$  are linearly independent in  $C^\infty(\mathbb{R})$ .

Suppose that  $ae^x + be^{2x} + ce^{3x} = 0$  for all  $x \in \mathbb{R}$ , where  $a, b, c$  are constants. We have to show that  $a = b = c = 0$ .

Differentiate this identity twice:

$$\begin{aligned}ae^x + be^{2x} + ce^{3x} &= 0, \\ae^x + 2be^{2x} + 3ce^{3x} &= 0, \\ae^x + 4be^{2x} + 9ce^{3x} &= 0.\end{aligned}$$

It follows that  $A(x)\mathbf{v} = \mathbf{0}$ , where

$$A(x) = \begin{pmatrix} e^x & e^{2x} & e^{3x} \\ e^x & 2e^{2x} & 3e^{3x} \\ e^x & 4e^{2x} & 9e^{3x} \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}.$$

$$A(x) = \begin{pmatrix} e^x & e^{2x} & e^{3x} \\ e^x & 2e^{2x} & 3e^{3x} \\ e^x & 4e^{2x} & 9e^{3x} \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}.$$

$$\begin{aligned} \det A(x) &= e^x \begin{vmatrix} 1 & e^{2x} & e^{3x} \\ 1 & 2e^{2x} & 3e^{3x} \\ 1 & 4e^{2x} & 9e^{3x} \end{vmatrix} = e^x e^{2x} \begin{vmatrix} 1 & 1 & e^{3x} \\ 1 & 2 & 3e^{3x} \\ 1 & 4 & 9e^{3x} \end{vmatrix} \\ &= e^x e^{2x} e^{3x} \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 4 & 9 \end{vmatrix} = e^{6x} \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 4 & 9 \end{vmatrix} = e^{6x} \begin{vmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 1 & 4 & 9 \end{vmatrix} \\ &= e^{6x} \begin{vmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 3 & 8 \end{vmatrix} = e^{6x} \begin{vmatrix} 1 & 2 \\ 3 & 8 \end{vmatrix} = 2e^{6x} \neq 0. \end{aligned}$$

Since the matrix  $A(x)$  is invertible, we obtain

$$A(x)\mathbf{v} = \mathbf{0} \implies \mathbf{v} = \mathbf{0} \implies a = b = c = 0$$

## Wronskian

Let  $f_1, f_2, \dots, f_n$  be smooth functions on an interval  $[a, b]$ . The **Wronskian**  $W[f_1, f_2, \dots, f_n]$  is a function on  $[a, b]$  defined by

$$W[f_1, f_2, \dots, f_n](x) = \begin{vmatrix} f_1(x) & f_2(x) & \cdots & f_n(x) \\ f_1'(x) & f_2'(x) & \cdots & f_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ f_1^{(n-1)}(x) & f_2^{(n-1)}(x) & \cdots & f_n^{(n-1)}(x) \end{vmatrix}.$$

**Theorem** If  $W[f_1, f_2, \dots, f_n](x_0) \neq 0$  for some  $x_0 \in [a, b]$  then the functions  $f_1, f_2, \dots, f_n$  are linearly independent in  $C[a, b]$ .

## Basis

*Definition.* Let  $V$  be a vector space. Any linearly independent spanning set for  $V$  is called a **basis**.

Suppose that a set  $S \subset V$  is a basis for  $V$ .

“Spanning set” means that any vector  $\mathbf{v} \in V$  can be represented as a linear combination

$$\mathbf{v} = r_1\mathbf{v}_1 + r_2\mathbf{v}_2 + \cdots + r_k\mathbf{v}_k,$$

where  $\mathbf{v}_1, \dots, \mathbf{v}_k$  are distinct vectors from  $S$  and  $r_1, \dots, r_k \in \mathbb{R}$ . “Linearly independent” implies that the above representation is unique:

$$\mathbf{v} = r_1\mathbf{v}_1 + r_2\mathbf{v}_2 + \cdots + r_k\mathbf{v}_k = r'_1\mathbf{v}_1 + r'_2\mathbf{v}_2 + \cdots + r'_k\mathbf{v}_k$$

$$\implies (r_1 - r'_1)\mathbf{v}_1 + (r_2 - r'_2)\mathbf{v}_2 + \cdots + (r_k - r'_k)\mathbf{v}_k = \mathbf{0}$$

$$\implies r_1 - r'_1 = r_2 - r'_2 = \cdots = r_k - r'_k = 0$$



## Basis

*Definition.* Let  $V$  be a vector space. Any linearly independent spanning set for  $V$  is called a **basis**.

**Theorem** A nonempty set  $S \subset V$  is a basis for  $V$  if and only if any vector  $\mathbf{v} \in V$  is *uniquely represented* as a linear combination

$\mathbf{v} = r_1\mathbf{v}_1 + r_2\mathbf{v}_2 + \cdots + r_k\mathbf{v}_k$ , where  $\mathbf{v}_1, \dots, \mathbf{v}_k$  are distinct vectors from  $S$  and  $r_1, \dots, r_k \in \mathbb{R}$ .

*Remark on uniqueness.* Expansions  $\mathbf{v} = 2\mathbf{v}_1 - \mathbf{v}_2$ ,  $\mathbf{v} = -\mathbf{v}_2 + 2\mathbf{v}_1$ , and  $\mathbf{v} = 2\mathbf{v}_1 - \mathbf{v}_2 + 0\mathbf{v}_3$  are considered the same.

*Examples.* • Standard basis for  $\mathbb{R}^n$ :

$$\mathbf{e}_1 = (1, 0, 0, \dots, 0, 0), \quad \mathbf{e}_2 = (0, 1, 0, \dots, 0, 0), \dots, \\ \mathbf{e}_n = (0, 0, 0, \dots, 0, 1).$$

Indeed,  $(x_1, x_2, \dots, x_n) = x_1\mathbf{e}_1 + x_2\mathbf{e}_2 + \dots + x_n\mathbf{e}_n$ .

- Matrices  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$

form a basis for  $\mathcal{M}_{2,2}(\mathbb{R})$ .

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + b \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + c \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + d \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

- Polynomials  $1, x, x^2, \dots, x^{n-1}$  form a basis for  $\mathcal{P}_n = \{a_0 + a_1x + \dots + a_{n-1}x^{n-1} : a_i \in \mathbb{R}\}$ .

- The infinite set  $\{1, x, x^2, \dots, x^n, \dots\}$  is a basis for  $\mathcal{P}$ , the space of all polynomials.

Let  $\mathbf{v}, \mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$  and  $r_1, r_2, \dots, r_k \in \mathbb{R}$ .

The vector equation  $r_1\mathbf{v}_1 + r_2\mathbf{v}_2 + \dots + r_k\mathbf{v}_k = \mathbf{v}$  is equivalent to the matrix equation  $A\mathbf{x} = \mathbf{v}$ , where

$$A = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k), \quad \mathbf{x} = \begin{pmatrix} r_1 \\ \vdots \\ r_k \end{pmatrix}.$$

---

$$r_1 \begin{pmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{n1} \end{pmatrix} + r_2 \begin{pmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{n2} \end{pmatrix} + \dots + r_k \begin{pmatrix} a_{1k} \\ a_{2k} \\ \vdots \\ a_{nk} \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix} \iff$$


$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1k} \\ a_{21} & a_{22} & \dots & a_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nk} \end{pmatrix} \begin{pmatrix} r_1 \\ r_2 \\ \vdots \\ r_k \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix} \iff A\mathbf{x} = \mathbf{v}$$

Let  $\mathbf{v}, \mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$  and  $r_1, r_2, \dots, r_k \in \mathbb{R}$ .  
The vector equation  $r_1\mathbf{v}_1 + r_2\mathbf{v}_2 + \dots + r_k\mathbf{v}_k = \mathbf{v}$  is  
equivalent to the matrix equation  $A\mathbf{x} = \mathbf{v}$ , where


$$A = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k), \quad \mathbf{x} = \begin{pmatrix} r_1 \\ \vdots \\ r_k \end{pmatrix}.$$

That is,  $A$  is the  $n \times k$  matrix such that vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  are consecutive columns of  $A$ .


- *Vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k$  span  $\mathbb{R}^n$  if the row echelon form of  $A$  has no zero rows.*
- *Vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k$  are linearly independent if the row echelon form of  $A$  has a leading entry in each column (no free variables).*




spanning  
linear independence



no spanning  
linear independence



spanning  
no linear independence



no spanning  
no linear independence

## Bases for $\mathbb{R}^n$

Let  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  be vectors in  $\mathbb{R}^n$ .

**Theorem 1** If  $k < n$  then the vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  do not span  $\mathbb{R}^n$ .

**Theorem 2** If  $k > n$  then the vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  are linearly dependent.

**Theorem 3** If  $k = n$  then the following conditions are equivalent:

- (i)  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  is a basis for  $\mathbb{R}^n$ ;
- (ii)  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  is a spanning set for  $\mathbb{R}^n$ ;
- (iii)  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  is a linearly independent set.

*Example.* Consider vectors  $\mathbf{v}_1 = (1, -1, 1)$ ,  $\mathbf{v}_2 = (1, 0, 0)$ ,  $\mathbf{v}_3 = (1, 1, 1)$ , and  $\mathbf{v}_4 = (1, 2, 4)$  in  $\mathbb{R}^3$ .

Vectors  $\mathbf{v}_1$  and  $\mathbf{v}_2$  are linearly independent (as they are not parallel), but they do not span  $\mathbb{R}^3$ .

Vectors  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  are linearly independent since

$$\begin{vmatrix} 1 & 1 & 1 \\ -1 & 0 & 1 \\ 1 & 0 & 1 \end{vmatrix} = - \begin{vmatrix} -1 & 1 \\ 1 & 1 \end{vmatrix} = -(-2) = 2 \neq 0.$$

Therefore  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is a basis for  $\mathbb{R}^3$ .

Vectors  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4$  span  $\mathbb{R}^3$  (because  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  already span  $\mathbb{R}^3$ ), but they are linearly dependent.

## Dimension

**Theorem 1** Any vector space has a basis.

**Theorem 2** If a vector space  $V$  has a finite basis, then all bases for  $V$  are finite and have the same number of elements.

*Definition.* The **dimension** of a vector space  $V$ , denoted  $\dim V$ , is the number of elements in any of its bases.



*Examples.* •  $\dim \mathbb{R}^n = n$

•  $\mathcal{M}_{2,2}(\mathbb{R})$ : the space of  $2 \times 2$  matrices  
 $\dim \mathcal{M}_{2,2}(\mathbb{R}) = 4$

•  $\mathcal{M}_{m,n}(\mathbb{R})$ : the space of  $m \times n$  matrices  
 $\dim \mathcal{M}_{m,n}(\mathbb{R}) = mn$

•  $\mathcal{P}_n$ : polynomials of degree less than  $n$   
 $\dim \mathcal{P}_n = n$

•  $\mathcal{P}$ : the space of all polynomials  
 $\dim \mathcal{P} = \infty$

•  $\{\mathbf{0}\}$ : the trivial vector space  
 $\dim \{\mathbf{0}\} = 0$