

MATH 415
Modern Algebra I

Lecture 23:
Some examples of rings.

Rings

Definition. A **ring** is a set R , together with two binary operations usually called **addition** and **multiplication** and denoted accordingly, such that

- R is an abelian group under addition,
- R is a semigroup under multiplication,
- multiplication distributes over addition.

The complete list of axioms is as follows:

(A0) for all $x, y \in R$, $x + y$ is an element of R ;

(A1) $(x + y) + z = x + (y + z)$ for all $x, y, z \in R$;

(A2) there exists an element, denoted 0 , in R such that $x + 0 = 0 + x = x$ for all $x \in R$;

(A3) for every $x \in R$ there exists an element, denoted $-x$, in R such that $x + (-x) = (-x) + x = 0$;

(A4) $x + y = y + x$ for all $x, y \in R$;

(M0) for all $x, y \in R$, xy is an element of R ;

(M1) $(xy)z = x(yz)$ for all $x, y, z \in R$;

(D) $x(y+z) = xy+xz$ and $(y+z)x = yx+zx$ for all $x, y, z \in R$.

Ring of functions

Let R be a ring and S be a nonempty set. Denote by $\mathcal{F}(S, R)$ the set of all functions $f : S \rightarrow R$. Given $f, g \in \mathcal{F}(S, R)$, we let $(f + g)(x) = f(x) + g(x)$ and $(fg)(x) = f(x)g(x)$ for all $x \in S$. That is, to add (or multiply) functions, we add (or multiply) their values at every point. Then $\mathcal{F}(S, R)$ is a ring.

The ring $\mathcal{F}(S, R)$ inherits many properties from the ring R , with one important exception. If R is a nontrivial ring and S has more than one element, then the ring $\mathcal{F}(S, R)$ has divisors of zero. Indeed, take any point $x_0 \in S$, any nonzero element $r \in R$, and let

$$f_1(x) = \begin{cases} r & \text{if } x = x_0, \\ 0 & \text{if } x \in S \setminus \{x_0\}; \end{cases} \quad f_2(x) = \begin{cases} 0 & \text{if } x = x_0, \\ r & \text{if } x \in S \setminus \{x_0\}. \end{cases}$$

Then the functions f_1 and f_2 are nonzero elements of the ring $\mathcal{F}(S, R)$ while $f_1 f_2 = 0$.

Ring of matrices

Let R be a ring. For any integers $m, n > 0$, denote by $\mathcal{M}_{m,n}(R)$ the set of all $m \times n$ matrices with entries from R . Given two matrices $A = (a_{ij})$ and $B = (b_{ij})$ in $\mathcal{M}_{m,n}(R)$, we let $A + B = (c_{ij})$ and $A - B = (d_{ij})$, where $c_{ij} = a_{ij} + b_{ij}$ and $d_{ij} = a_{ij} - b_{ij}$, $1 \leq i \leq m$, $1 \leq j \leq n$. Given matrices $A = (a_{ij}) \in \mathcal{M}_{m,n}(R)$ and $B = (b_{ij}) \in \mathcal{M}_{n,p}(R)$, we let $AB = (c_{ij})$, where $c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}$, $1 \leq i \leq m$, $1 \leq j \leq p$.

Matrix multiplication is associative. Indeed, let $A = (a_{ij}) \in \mathcal{M}_{m,n}(R)$, $B = (b_{jk}) \in \mathcal{M}_{n,p}(R)$ and $C = (c_{kl}) \in \mathcal{M}_{p,q}(R)$. Then $(AB)C = (d_{i\ell})$ and $A(BC) = (d'_{i\ell})$ are matrices in $\mathcal{M}_{m,q}(R)$. Using distributive laws in R , we obtain that

$$d_{i\ell} = \sum_{k=1}^p \sum_{j=1}^n (a_{ij} b_{jk}) c_{k\ell}, \quad d'_{i\ell} = \sum_{j=1}^n \sum_{k=1}^p a_{ij} (b_{jk} c_{k\ell}).$$

Hence $(AB)C = A(BC)$ since R is a ring.

As a consequence, square matrices in $\mathcal{M}_{n,n}(R)$ form a ring.

Direct product of rings

Suppose R_1, R_2, \dots, R_n are rings. We define addition and multiplication on the Cartesian product $R_1 \times R_2 \times \dots \times R_n$ by

$$(r_1, r_2, \dots, r_n) + (r'_1, r'_2, \dots, r'_n) = (r_1 + r'_1, r_2 + r'_2, \dots, r_n + r'_n),$$
$$(r_1, r_2, \dots, r_n)(r'_1, r'_2, \dots, r'_n) = (r_1 r'_1, r_2 r'_2, \dots, r_n r'_n)$$

for all $r_i, r'_i \in R_i$, $1 \leq i \leq n$.

Then $R_1 \times R_2 \times \dots \times R_n$ is a ring called the **direct product** of rings R_1, R_2, \dots, R_n .

The ring $R_1 \times R_2 \times \dots \times R_n$ is commutative if each of the rings R_1, R_2, \dots, R_n is commutative. It is a ring with unity if each of the rings R_1, R_2, \dots, R_n has the unity.

If at least two of the rings R_1, R_2, \dots, R_n are nontrivial, then the direct product $R_1 \times R_2 \times \dots \times R_n$ admits divisors of zero.

Complex numbers

\mathbb{C} : complex numbers.

Complex number: $z = x + iy,$

where $x, y \in \mathbb{R}$ and $i^2 = -1$.

$i = \sqrt{-1}$: imaginary unit

Alternative notation: $z = x + yi$.

x = real part of z ,

iy = imaginary part of z

$y = 0 \implies z = x$ (real number)

$x = 0 \implies z = iy$ (purely imaginary number)

We add, subtract, and multiply complex numbers as polynomials in i (but keep in mind that $i^2 = -1$).

If $z_1 = x_1 + iy_1$ and $z_2 = x_2 + iy_2$, then

$$z_1 + z_2 = (x_1 + x_2) + i(y_1 + y_2),$$

$$z_1 - z_2 = (x_1 - x_2) + i(y_1 - y_2),$$

$$z_1 z_2 = (x_1 x_2 - y_1 y_2) + i(x_1 y_2 + x_2 y_1).$$

Given $z = x + iy$, the **complex conjugate** of z is $\bar{z} = x - iy$. The **modulus** of z is $|z| = \sqrt{x^2 + y^2}$.

$$z\bar{z} = (x + iy)(x - iy) = x^2 - (iy)^2 = x^2 + y^2 = |z|^2.$$

$$z^{-1} = \frac{\bar{z}}{|z|^2}, \quad (x + iy)^{-1} = \frac{x - iy}{x^2 + y^2}.$$

Complex exponentials

Definition. For any $z \in \mathbb{C}$ let

$$e^z = 1 + z + \frac{z^2}{2!} + \cdots + \frac{z^n}{n!} + \cdots$$

Remark. A sequence of complex numbers $z_1 = x_1 + iy_1$, $z_2 = x_2 + iy_2$, \dots converges to $z = x + iy$ if $x_n \rightarrow x$ and $y_n \rightarrow y$ as $n \rightarrow \infty$.

Theorem 1 If $z = x + iy$, $x, y \in \mathbb{R}$, then

$$e^z = e^x(\cos y + i \sin y).$$

In particular, $e^{i\phi} = \cos \phi + i \sin \phi$, $\phi \in \mathbb{R}$.

Theorem 2 $e^{z+w} = e^z \cdot e^w$ for all $z, w \in \mathbb{C}$.

Proposition $e^{i\phi} = \cos \phi + i \sin \phi$ for all $\phi \in \mathbb{R}$.

Proof:
$$e^{i\phi} = 1 + i\phi + \frac{(i\phi)^2}{2!} + \dots + \frac{(i\phi)^n}{n!} + \dots$$

The sequence $1, i, i^2, i^3, \dots, i^n, \dots$ is periodic:



$$\underbrace{1, i, -1, -i}, \underbrace{1, i, -1, -i}, \dots$$

It follows that

$$\begin{aligned} e^{i\phi} &= 1 - \frac{\phi^2}{2!} + \frac{\phi^4}{4!} - \dots + (-1)^k \frac{\phi^{2k}}{(2k)!} + \dots \\ &+ i \left(\phi - \frac{\phi^3}{3!} + \frac{\phi^5}{5!} - \dots + (-1)^k \frac{\phi^{2k+1}}{(2k+1)!} + \dots \right) \\ &= \cos \phi + i \sin \phi. \end{aligned}$$

Geometric representation

Any complex number $z = x + iy$ is represented by the vector/point $(x, y) \in \mathbb{R}^2$.



$$x = r \cos \phi, \quad y = r \sin \phi \implies z = r(\cos \phi + i \sin \phi) = re^{i\phi}$$

If $z_1 = r_1 e^{i\phi_1}$ and $z_2 = r_2 e^{i\phi_2}$, then

$$z_1 z_2 = r_1 r_2 e^{i(\phi_1 + \phi_2)}, \quad z_1 / z_2 = (r_1 / r_2) e^{i(\phi_1 - \phi_2)}.$$

Complex numbers as an \mathbb{R} -algebra

Complex numbers can be defined as a certain 2-dimensional algebra over the field \mathbb{R} . We have a distinguished basis $\mathbf{1}, i$. Hence every complex number z is uniquely represented as $z = x\mathbf{1} + yi$, where $x, y \in \mathbb{R}$.

Since multiplication is a bilinear function, it is enough to define $z_1 \cdot z_2$ in the case $z_1, z_2 \in \{\mathbf{1}, i\}$. We set $\mathbf{1} \cdot \mathbf{1} = \mathbf{1}$, $\mathbf{1} \cdot i = i \cdot \mathbf{1} = i$ and $i \cdot i = -\mathbf{1}$.

Because of bilinearity of the product, it is easy to check that $\mathbf{1} \cdot z = z \cdot \mathbf{1}$, $z_1 \cdot z_2 = z_2 \cdot z_1$ and $(z_1 \cdot z_2) \cdot z_3 = z_1 \cdot (z_2 \cdot z_3)$.