

MATH 433  
Applied Algebra

**Lecture 14:**  
**Functions.**  
**Relations.**

## Cartesian product

*Definition.* The **Cartesian product**  $X \times Y$  of two sets  $X$  and  $Y$  is the set of all ordered pairs  $(x, y)$  such that  $x \in X$  and  $y \in Y$ .

The Cartesian square  $X \times X$  is also denoted  $X^2$ .

If the sets  $X$  and  $Y$  are finite, then

$\#(X \times Y) = (\#X)(\#Y)$ , where  $\#S$  denote the number of elements in a set  $S$ .

## Functions


A **function**  $f : X \rightarrow Y$  is an assignment: to each  $x \in X$  we assign an element  $f(x) \in Y$ .

The **graph** of the function  $f : X \rightarrow Y$  is defined as the subset of  $X \times Y$  consisting of all pairs of the form  $(x, f(x))$ ,  $x \in X$ .

*Definition.* A function  $f : X \rightarrow Y$  is **surjective** (or **onto**) if for each  $y \in Y$  there exists at least one  $x \in X$  such that  $f(x) = y$ . The function  $f$  is **injective** (or **one-to-one**) if  $f(x') = f(x) \implies x' = x$ .

Finally,  $f$  is **bijective** if it is both surjective and injective. Equivalently, if for each  $y \in Y$  there is exactly one  $x \in X$  such that  $f(x) = y$ .

A function  $g : Y \rightarrow X$  is called the **inverse** of  $f$  (and denoted  $f^{-1}$ ) if  $y = f(x) \iff x = g(y)$  for all  $x \in X$  and  $y \in Y$ . The inverse function  $f^{-1}$  exists if and only if  $f$  is bijective.



## Relations

*Definition.* Let  $X$  and  $Y$  be sets. A **relation**  $R$  from  $X$  to  $Y$  is given by specifying a subset of the Cartesian product:  $S_R \subset X \times Y$ .

If  $(x, y) \in S_R$ , then we say that  $x$  **is related to**  $y$  (in the sense of  $R$  or by  $R$ ) and write  $xRy$ .

*Remarks.* • Usually the relation  $R$  is identified with the set  $S_R$ .

• In the case  $X = Y$ , the relation  $R$  is called a **relation on**  $X$ .

**Examples.** • “is equal to”

$$xRy \iff x = y$$

Equivalently,  $R = \{(x, x) \mid x \in X \cap Y\}$ .

• “is not equal to”

$$xRy \iff x \neq y$$

• “is mapped by  $f$  to”

$xRy \iff y = f(x)$ , where  $f : X \rightarrow Y$  is a function.

Equivalently,  $R$  is the graph of the function  $f$ .

• “is the image under  $f$  of”

(from  $Y$  to  $X$ )  $yRx \iff y = f(x)$ , where  $f : X \rightarrow Y$  is a function. If  $f$  is invertible, then  $R$  is the graph of  $f^{-1}$ .

• reversed  $R'$

$xRy \iff yR'x$ , where  $R'$  is a relation from  $Y$  to  $X$ .

• not  $R'$

$xRy \iff \text{not } xR'y$ , where  $R'$  is a relation from  $X$  to  $Y$ .

Equivalently,  $R = (X \times Y) \setminus R'$  (set difference).

## Relations on a set

- “is equal to”

$$xRy \iff x = y$$

- “is not equal to”

$$xRy \iff x \neq y$$

- “is less than”

$$X = \mathbb{R}, \quad xRy \iff x < y$$

- “is less than or equal to”

$$X = \mathbb{R}, \quad xRy \iff x \leq y$$

- “is contained in”

$X =$  the set of all subsets of some set  $Y$ ,

$$xRy \iff x \subset y$$

- “is congruent modulo  $n$  to”

$$X = \mathbb{Z}, \quad xRy \iff x \equiv y \pmod{n}$$

- “divides”

$$X = \mathbb{P}, \quad xRy \iff x|y$$


A relation  $R$  on a finite set  $X$  can be represented by a **directed graph**.

Vertices of the graph are elements of  $X$ , and we have a directed edge from  $x$  to  $y$  if and only if  $xRy$ .


Another way to represent the relation  $R$  is the **adjacency table**.

Rows and columns are labeled by elements of  $X$ . We put 1 at the intersection of a row  $x$  with a column  $y$  if  $xRy$ . Otherwise we put 0.





	$a$	$b$	$c$
$a$	0	1	1
$b$	0	1	1
$c$	1	0	0



	$a$	$b$	$c$	$d$	$e$
$a$	0	1	1	0	0
$b$	0	1	1	0	0
$c$	0	0	0	1	0
$d$	0	0	0	0	1
$e$	1	0	0	0	0

## Properties of relations

*Definition.* Let  $R$  be a relation on a set  $X$ . We say that  $R$  is

- **reflexive** if  $xRx$  for all  $x \in X$ ,
- **symmetric** if, for all  $x, y \in X$ ,  $xRy$  implies  $yRx$ ,
- **antisymmetric** if, for all  $x, y \in X$ ,  $xRy$  and  $yRx$  cannot hold simultaneously,
- **weakly antisymmetric** if, for all  $x, y \in X$ ,  $xRy$  and  $yRx$  imply that  $x = y$ ,
- **transitive** if, for all  $x, y, z \in X$ ,  $xRy$  and  $yRz$  imply that  $xRz$ .

## Partial ordering

*Definition.* A relation  $R$  on a set  $X$  is a **partial ordering** (or **partial order**) if  $R$  is reflexive, weakly antisymmetric, and transitive:

- $xRx$ ,
- $xRy$  and  $yRx \implies x = y$ ,
- $xRy$  and  $yRz \implies xRz$ .

A relation  $R$  on a set  $X$  is a **strict partial order** if  $R$  is antisymmetric and transitive:

- $xRy \implies \text{not } yRx$ ,
- $xRy$  and  $yRz \implies xRz$ .

*Examples.* “is less than or equal to”, “is contained in”, “is a divisor of” are partial orders. “is less than” is a strict order.

## Equivalence relation

*Definition.* A relation  $R$  on a set  $X$  is an **equivalence relation** if  $R$  is reflexive, symmetric, and transitive:

- $xRx$ ,
- $xRy \implies yRx$ ,
- $xRy$  and  $yRz \implies xRz$ .

*Examples.* “is equal to”, “is congruent modulo  $n$  to” are equivalence relations.

Given an equivalence relation  $R$  on  $X$ , the **equivalence class** of an element  $x \in X$  relative to  $R$  is the set of all elements  $y \in X$  such that  $yRx$ .

**Theorem** The equivalence classes form a **partition** of the set  $X$ , which means that

- any two equivalence classes either coincide, or else they are disjoint,
- any element of  $X$  belongs to some equivalence class.