

MATH 433
Applied Algebra

Lecture 19:
Sign of a permutation.
Classical definition of the determinant.

Sign of a permutation

Theorem 1 (i) Any permutation of $n \geq 2$ elements is a product of transpositions. (ii) If $\pi = \tau_1 \tau_2 \dots \tau_k = \tau'_1 \tau'_2 \dots \tau'_m$, where τ_i, τ'_j are transpositions, then the numbers k and m are of the same parity (that is, both even or both odd).

A permutation π is called **even** if it is a product of an even number of transpositions, and **odd** if it is a product of an odd number of transpositions.

The **sign** $\text{sgn}(\pi)$ of the permutation π is defined to be $+1$ if π is even, and -1 if π is odd.

Theorem 2 (i) $\text{sgn}(\pi\sigma) = \text{sgn}(\pi)\text{sgn}(\sigma)$ for any $\pi, \sigma \in S_X$.

(ii) $\text{sgn}(\pi^{-1}) = \text{sgn}(\pi)$ for any $\pi \in S_X$.

(iii) $\text{sgn}(\text{id}) = 1$.

(iv) $\text{sgn}(\tau) = -1$ for any transposition τ .

(v) $\text{sgn}(\sigma) = (-1)^{r-1}$ for any cycle σ of length r .

Examples

- $\pi = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 \\ 2 & 4 & 7 & 9 & 1 & 12 & 5 & 11 & 3 & 10 & 6 & 8 \end{pmatrix}.$

First we decompose π into a product of disjoint cycles:

$$\pi = (1\ 2\ 4\ 9\ 3\ 7\ 5)(6\ 12\ 8\ 11).$$

The cycle $\sigma_1 = (1\ 2\ 4\ 9\ 3\ 7\ 5)$ has length 7, hence it is an even permutation. The cycle $\sigma_2 = (6\ 12\ 8\ 11)$ has length 4, hence it is an odd permutation. Then

$$\text{sgn}(\pi) = \text{sgn}(\sigma_1\sigma_2) = \text{sgn}(\sigma_1)\text{sgn}(\sigma_2) = 1 \cdot (-1) = -1.$$

- $\pi = (2\ 4\ 3)(1\ 2)(2\ 3\ 4).$

π is represented as a product of cycles. The transposition has sign -1 while the cycles of length 3 have sign $+1$. Even though the cycles are not disjoint, $\text{sgn}(\pi) = 1 \cdot (-1) \cdot 1 = -1$.

Let $\pi \in S(n)$ and i, j be integers, $1 \leq i < j \leq n$. We say that the permutation π preserves order of the pair (i, j) if $\pi(i) < \pi(j)$. Otherwise π makes an **inversion**. Denote by $N(\pi)$ the number of inversions made by the permutation π .

Lemma 1 Let $\tau, \pi \in S(n)$ and suppose that τ is an adjacent transposition, $\tau = (k \ k+1)$. Then $|N(\tau\pi) - N(\pi)| = 1$.

Proof: For every pair (i, j) , $1 \leq i < j \leq n$, let us compare the order of pairs $\pi(i), \pi(j)$ and $\tau\pi(i), \tau\pi(j)$. We observe that the order differs exactly for one pair, when $\{\pi(i), \pi(j)\} = \{k, k+1\}$. The lemma follows.

Lemma 2 Let $\pi \in S(n)$ and $\tau_1, \tau_2, \dots, \tau_k$ be adjacent transpositions. Then **(i)** for any $\pi \in S(n)$ the numbers k and $N(\tau_1\tau_2 \dots \tau_k\pi) - N(\pi)$ are of the same parity, **(ii)** the numbers k and $N(\tau_1\tau_2 \dots \tau_k)$ are of the same parity.

Sketch of the proof: **(i)** follows from Lemma 1 by induction on k . **(ii)** is a particular case of part (i), when $\pi = \text{id}$.

Lemma 3 (i) Any cycle of length r is a product of $r-1$ transpositions. **(ii)** Any transposition is a product of an odd number of adjacent transpositions.

Proof: **(i)** $(x_1 x_2 \dots x_r) = (x_1 x_2)(x_2 x_3)(x_3 x_4) \dots (x_{r-1} x_r)$.

(ii) $(k k+r) = \sigma^{-1}(k k+1)\sigma$, where $\sigma = (k+1 k+2 \dots k+r)$.

By the above, $\sigma = (k+1 k+2)(k+2 k+3) \dots (k+r-1 k+r)$
and $\sigma^{-1} = (k+r k+r-1) \dots (k+3 k+2)(k+2 k+1)$.

Theorem (i) Any permutation is a product of transpositions.

(ii) If $\pi = \tau_1 \tau_2 \dots \tau_k$, where τ_i are transpositions, then the numbers k and $N(\pi)$ are of the same parity.

Proof: **(i)** Any permutation is a product of disjoint cycles.

By Lemma 3, any cycle is a product of transpositions.

(ii) By Lemma 3, each of $\tau_1, \tau_2, \dots, \tau_k$ is a product of an odd number of adjacent transpositions. Hence $\pi = \tau'_1 \tau'_2 \dots \tau'_m$, where τ'_i are adjacent transpositions and number m is of the same parity as k . By Lemma 2, m has the same parity as $N(\pi)$.

Alternating group

Given an integer $n \geq 2$, the **alternating group** on n symbols, denoted A_n or $A(n)$, is the set of all even permutations in the symmetric group $S(n)$.

Theorem (i) For any two permutations $\pi, \sigma \in A(n)$, the product $\pi\sigma$ is also in $A(n)$.

(ii) The identity function id is in $A(n)$.

(iii) For any permutation $\pi \in A(n)$, the inverse π^{-1} is in $A(n)$.

Theorem The alternating group $A(n)$ has $n!/2$ elements.

Proof: Consider a function $F : S(n) \rightarrow S(n)$ given by $F(\pi) = (1\ 2)\pi$. Note that F is bijective (indeed, $F^{-1} = F$). Hence $|F(E)| = |E|$ for any set $E \subset S(n)$. We observe that $F(A(n)) \subset S(n) \setminus A(n)$ and $F(S(n) \setminus A(n)) \subset A(n)$. Therefore $|A(n)| \leq |S(n) \setminus A(n)|$ and $|S(n) \setminus A(n)| \leq |A(n)|$ so that $|A(n)| = |S(n) \setminus A(n)| = |S(n)|/2 = n!/2$.

Examples. • The alternating group $A(3)$ has 3 elements: the identity function and two cycles of length 3, $(1\ 2\ 3)$ and $(1\ 3\ 2)$.

• The alternating group $A(4)$ has 12 elements of the following **cycle shapes**: id, $(1\ 2\ 3)$, and $(1\ 2)(3\ 4)$.

• The alternating group $A(5)$ has 60 elements of the following cycle shapes: id, $(1\ 2\ 3)$, $(1\ 2)(3\ 4)$, and $(1\ 2\ 3\ 4\ 5)$.

Classical definition of the determinant

Definition. $\det(a) = a$, $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$,

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} \\ - a_{13}a_{22}a_{31} - a_{12}a_{21}a_{33} - a_{11}a_{23}a_{32}.$$

If $A = (a_{ij})$ is an $n \times n$ matrix then

$$\det A = \sum_{\pi \in S(n)} \operatorname{sgn}(\pi) a_{1,\pi(1)} a_{2,\pi(2)} \cdots a_{n,\pi(n)},$$

where π runs over all permutations of $\{1, 2, \dots, n\}$.

Theorem $\det A^T = \det A$.

Proof: Let $A = (a_{ij})_{1 \leq i, j \leq n}$. Then $A^T = (b_{ij})_{1 \leq i, j \leq n}$, where $b_{ij} = a_{ji}$. We have

$$\begin{aligned}\det A^T &= \sum_{\pi \in S(n)} \operatorname{sgn}(\pi) b_{1, \pi(1)} b_{2, \pi(2)} \cdots b_{n, \pi(n)} \\ &= \sum_{\pi \in S(n)} \operatorname{sgn}(\pi) a_{\pi(1), 1} a_{\pi(2), 2} \cdots a_{\pi(n), n} \\ &= \sum_{\pi \in S(n)} \operatorname{sgn}(\pi) a_{1, \pi^{-1}(1)} a_{2, \pi^{-1}(2)} \cdots a_{n, \pi^{-1}(n)}.\end{aligned}$$

When π runs over all permutations of $\{1, 2, \dots, n\}$, so does $\sigma = \pi^{-1}$. It follows that

$$\begin{aligned}\det A^T &= \sum_{\sigma \in S(n)} \operatorname{sgn}(\sigma^{-1}) a_{1, \sigma(1)} a_{2, \sigma(2)} \cdots a_{n, \sigma(n)} \\ &= \sum_{\sigma \in S(n)} \operatorname{sgn}(\sigma) a_{1, \sigma(1)} a_{2, \sigma(2)} \cdots a_{n, \sigma(n)} = \det A.\end{aligned}$$

Theorem 1 Suppose A is a square matrix and B is obtained from A by exchanging two rows. Then $\det B = -\det A$.

Theorem 2 Suppose A is a square matrix and B is obtained from A by permuting its rows. Then $\det B = \det A$ if the permutation is even and $\det B = -\det A$ if the permutation is odd.

Proof: Let $A = (a_{ij})_{1 \leq i, j \leq n}$ be an $n \times n$ matrix. Suppose that a matrix B is obtained from A by permuting its rows according to a permutation $\sigma \in S(n)$. Then $B = (b_{ij})_{1 \leq i, j \leq n}$, where $b_{\sigma(i), j} = a_{ij}$. Equivalently, $b_{ij} = a_{\sigma^{-1}(i), j}$. We have

$$\begin{aligned} \det B &= \sum_{\pi \in S(n)} \operatorname{sgn}(\pi) b_{1, \pi(1)} b_{2, \pi(2)} \cdots b_{n, \pi(n)} \\ &= \sum_{\pi \in S(n)} \operatorname{sgn}(\pi) a_{\sigma^{-1}(1), \pi(1)} a_{\sigma^{-1}(2), \pi(2)} \cdots a_{\sigma^{-1}(n), \pi(n)} \\ &= \sum_{\pi \in S(n)} \operatorname{sgn}(\pi) a_{1, \pi\sigma(1)} a_{2, \pi\sigma(2)} \cdots a_{n, \pi\sigma(n)}. \end{aligned}$$

When π runs over all permutations of $\{1, 2, \dots, n\}$, so does $\tau = \pi\sigma$. It follows that

$$\begin{aligned} \det B &= \sum_{\tau \in S(n)} \operatorname{sgn}(\tau\sigma^{-1}) a_{1, \tau(1)} a_{2, \tau(2)} \cdots a_{n, \tau(n)} \\ &= \operatorname{sgn}(\sigma^{-1}) \sum_{\tau \in S(n)} \operatorname{sgn}(\tau) a_{1, \tau(1)} a_{2, \tau(2)} \cdots a_{n, \tau(n)} = \operatorname{sgn}(\sigma) \det A. \end{aligned}$$

The Vandermonde determinant

Definition. The **Vandermonde determinant** is the determinant of the following matrix

$$V = \begin{pmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{n-1} \\ 1 & x_3 & x_3^2 & \cdots & x_3^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_n & x_n^2 & \cdots & x_n^{n-1} \end{pmatrix},$$

where $x_1, x_2, \dots, x_n \in \mathbb{R}$. Equivalently, $V = (a_{ij})_{1 \leq i, j \leq n}$, where $a_{ij} = x_i^{j-1}$.

Theorem

$$\begin{vmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{n-1} \\ 1 & x_3 & x_3^2 & \cdots & x_3^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_n & x_n^2 & \cdots & x_n^{n-1} \end{vmatrix} = \prod_{1 \leq i < j \leq n} (x_j - x_i).$$

Corollary Consider a polynomial

$$p(x_1, x_2, \dots, x_n) = \prod_{1 \leq i < j \leq n} (x_j - x_i).$$

Then

$$p(x_{\pi(1)}, x_{\pi(2)}, \dots, x_{\pi(n)}) = \operatorname{sgn}(\pi) p(x_1, x_2, \dots, x_n)$$

for any permutation $\pi \in S(n)$.