

MATH 614

Dynamical Systems and Chaos

**Lecture 13:**  
**Bifurcation theory.**

## Bifurcation theory

The object of **bifurcation theory** is to study changes that maps undergo as parameters change.


In the context of one-dimensional dynamics, we consider a one-parameter family of maps  $f_\lambda : \mathbb{R} \rightarrow \mathbb{R}$ . We assume that  $G(x, \lambda) = f_\lambda(x)$  is smooth a function of two variables.

Informally, the family  $\{f_\lambda\}$  has a **bifurcation** at  $\lambda = \lambda_0$  if the dynamics of  $f_\lambda$  changes as  $\lambda$  passes  $\lambda_0$ . One way to formalize it is to require that there exist  $\varepsilon > 0$  such that for any  $\varepsilon_1, \varepsilon_2 \in (0, \varepsilon)$  the maps  $f_{\lambda_0 - \varepsilon_1}$  and  $f_{\lambda_0 + \varepsilon_2}$  are not topologically conjugate. The simplest case is an isolated bifurcation point  $\lambda_0$ . In this case, the map  $f_\lambda$  is structurally stable for all  $\lambda$  in a punctured neighborhood of  $\lambda_0$  but not for  $\lambda = \lambda_0$ .

The condition of topological conjugacy is often relaxed to local topological conjugacy or to similar configuration of periodic orbits.

## Saddle-node bifurcation


Exponential map  $E_\lambda(x) = \lambda e^x$ ,  $\lambda \approx 1/e$ ,  $x \approx 1$ .



For  $\lambda > 1/e$ , there are no fixed points. At  $\lambda = 1/e$ , there is a non-hyperbolic fixed point 1. For  $0 < \lambda < 1/e$ , there are two fixed points, one is repelling and the other one is attracting.


# Saddle-node bifurcation

Exponential map  $E_\lambda(x) = \lambda e^x$ ,  $\lambda \approx 1/e$ ,  $x \approx 1$ .




## Bifurcation diagram (saddle-node bifurcation)

In the plane with coordinates  $(\lambda, x)$ , we plot fixed points of  $E_\lambda$  for each  $\lambda$ :



## Period doubling bifurcation


Exponential map  $E_\lambda(x) = \lambda e^x$ ,  $\lambda \approx -e$ ,  $x \approx -1$ .



For  $-e < \lambda < 0$ , the fixed point is attracting. At  $\lambda = -e$ , it is not hyperbolic. For  $\lambda < -e$ , the fixed point is repelling and there is also an attracting periodic orbit of period 2.

## Bifurcation diagram (period doubling bifurcation)


In the plane with coordinates  $(\lambda, x)$ , we plot fixed points of  $E_\lambda^2$  for each  $\lambda$ :



## Period doubling: logistic map

Logistic map  $F_\mu(x) = \mu x(1-x)$ ,  $\mu \approx 3$ ,  $x \approx 2/3$ .

Consider graphs of  $F_\mu^2$  for  $\mu \approx 3$ :





For  $\mu < 3$ , the fixed point  $p_\mu = 1 - \mu^{-1}$  is attracting. At  $\mu = 3$ , it is not hyperbolic. For  $\mu > 3$ , the fixed point  $p_\mu$  is repelling and there is also an attracting periodic orbit of period 2.



## No bifurcation: sufficient condition

**Theorem 1** Let  $f_\lambda$  be a one-parameter family of functions and suppose that  $f_{\lambda_0}(x_0) = x_0$  and  $f'_{\lambda_0}(x_0) \neq 1$ . Then there are open intervals  $I \ni x_0$  and  $N \ni \lambda_0$  and a smooth function  $p : N \rightarrow I$  such that  $p(\lambda_0) = x_0$  and  $f_\lambda(p(\lambda)) = p(\lambda)$  for all  $\lambda \in N$ . Moreover,  $p(\lambda)$  is the only fixed point of  $f_\lambda$  in  $I$ .



## No bifurcation: sufficient condition

**Theorem 1** Let  $f_\lambda$  be a one-parameter family of functions and suppose that  $f_{\lambda_0}(x_0) = x_0$  and  $f'_{\lambda_0}(x_0) \neq 1$ . Then there are open intervals  $I \ni x_0$  and  $N \ni \lambda_0$  and a smooth function  $p : N \rightarrow I$  such that  $p(\lambda_0) = x_0$  and  $f_\lambda(p(\lambda)) = p(\lambda)$  for all  $\lambda \in N$ . Moreover,  $p(\lambda)$  is the only fixed point of  $f_\lambda$  in  $I$ .


*Proof:* Consider a function of two variables

$G(x, \lambda) = f_\lambda(x) - x$ . We have  $G(x_0, \lambda_0) = f_{\lambda_0}(x_0) - x_0 = 0$  and  $\frac{\partial G}{\partial x}(x_0, \lambda_0) = f'_{\lambda_0}(x_0) - 1 \neq 0$ . By the Implicit Function Theorem, there are open intervals  $I \ni x_0$  and  $N \ni \lambda_0$  and a smooth function  $p : N \rightarrow I$  such that

$$G(x, \lambda) = 0 \iff x = p(\lambda) \text{ for all } (x, \lambda) \in I \times N.$$


## Saddle-node bifurcation: sufficient condition

**Theorem 2** Let  $f_\lambda$  be a one-parameter family of functions and suppose that  $f_{\lambda_0}(x_0) = x_0$ ,  $f'_{\lambda_0}(x_0) = 1$ ,  $f''_{\lambda_0}(x_0) \neq 0$ , and  $\frac{\partial f_\lambda}{\partial \lambda} \Big|_{\lambda=\lambda_0}(x_0) \neq 0$ . Then there are open intervals  $I \ni x_0$  and  $N \ni \lambda_0$  and a smooth function  $p : I \rightarrow N$  such that  $p(x_0) = \lambda_0$  and  $f_{p(x)}(x) = x$  for all  $x \in I$ . Moreover,  $p'(x_0) = 0$  and  $p''(x_0) \neq 0$ .



## Period doubling bifurcation: sufficient condition

**Theorem 3** Let  $f_\lambda$  be a one-parameter family of functions and suppose that  $f_\lambda(x_0) = x_0$  for all  $\lambda$ ,  $f'_{\lambda_0}(x_0) = -1$ , and  $\frac{\partial(f_\lambda^2)'}{\partial\lambda} \Big|_{\lambda=\lambda_0}(x_0) \neq 0$ . Then there are open intervals  $I \ni x_0$  and  $N \ni \lambda_0$  and a smooth function  $p : I \rightarrow N$  such that  $p(x_0) = \lambda_0$  and  $f_{p(x)}^2(x) = x$  for all  $x \in I$  but  $f_{p(x)}(x) \neq x$  for  $x \in I \setminus \{x_0\}$ .



## More examples

- Quadratic maps:  $Q_c(x) = x^2 + c$ .

The family undergoes a saddle-node bifurcation at  $c = 1/4$  and a period doubling bifurcation at  $c = -3/4$ . It undergoes a lot of other bifurcations as well.

- Hyperbolic sine family:  $H_\lambda(x) = \lambda \sinh x$ .

A map  $H_\lambda$  is not structurally stable within the family for  $\lambda = -1, 0$ , and  $1$ . At  $\lambda = -1$ , we have a period doubling bifurcation. At  $\lambda = 1$ , the family transitions from one to three fixed points. At  $\lambda = 0$ , the bifurcation does not change the configuration of periodic points.

- Linear maps:  $f_\lambda(x) = \lambda^2 x$ .

A map  $f_\lambda$  is not structurally stable within the family for  $\lambda = -1, 0$ , and  $1$ . At  $\lambda = -1$  and  $1$ , the family transitions from a repelling fixed point to an attracting one (or vice versa). At  $\lambda = 0$ , there is no bifurcation.